Table 3.—Solar radiation measurements, and determinations of atmospheric factor,  $\beta$ , Washington, D. C., January, 1933—Con.

[Values in italics have been interpolated]

Date and solar hour angle	Solar alti- tude, h.	Air mass, m.	I,	I,	Ιr	β	Blue- ness of sky		Notes: Sky- light polari- zation, P., clouds, etc.
1933 Jan. 24 2:38 a	21-08 20-22 25-04 30-38 30-59 31-54 31-54 29-45 29-30 24-39 24-13	2. 76 2. 61 2. 36 1. 96 1. 89 1. 89 2. 01 2. 02 2. 39 2. 43	gr. cal. 1. 107 1. 139 1. 145 1. 241 1. 274 1. 276 1. 300 1. 292 1. 303 1. 200 1. 203	gr. cal, .853 .856 .870 .945 .945 .945 .945 .918 .918 .967	gr. cal. .693 .699 .703 .711 .714 .747 .748 .746 .747 .705 .708	. 070 . 070 . 080 . 070 . 065 . 075 . 060 . 055	4		P=55.8.
Jan. 30 3:28 a	15-28 16-31 23-31 24-15 28-56 29-21	3. 70 3. 49 2. 50 2. 43 2. 06 2. 04	1. 067 1. 150 1. 200 1. 216 1. 355 1. 310	. 964 . 967 . 948 . 951 . 999 1. 015	.702 .704 .786 .789 .814 .817	. 065 . 040 . 045 . 045 . 060 . 085		355	
Jan. 31 3:36 a 3:31 a 3:08 a 3:02 a 3:42 a 2:03 a 1:58 a 1:22 a 1:18 a	14-27 15-14 18-39 19-30 22-15 23-16 26-54 27-24 30-35 30-53	3. 97 3. 77 3. 11 2. 98 2. 63 2. 51 2. 21 2. 17 1. 96 1. 94	. 909 . 906 1. 016 1. 029 1. 032 1. 126 1. 174 1. 174 1. 221 1. 243	.717 .720 .802 .804 .846 .847 .885 .885 .897	.606 .609 .655 .658 .695 .697 .711 .712 .723	. 070 . 080 . 080 . 085 . 115 . 080 . 085 . 085 . 080 . 075	5	389	P=58. 3.

## POSITIONS AND AREAS OF SUN SPOTS

[Communicated by Capt. J. F. Hellweg, Superintendent United States Nava Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of longitude are measured from central meridian, positive west. The north latitudes are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column

	Eastern		Heliographic			Агеа		Total area
Date		standard civil time		Longi- tude	Lati- tude	Spot	Group	for each day
Jan. 1 (Naval Observatory)	13	m. 29 40 23				185	185	185
Jan. 5 (Naval Observatory)	14	6	-59. 0 -59. 0 -44. 0	328. 3 313. 5 328. 5	$+11.0 \\ +10.0 \\ +10.0$	170 154	154	355 308
Jan. 6 (Naval Observatory) Jan. 7 (Mount Wilson)	12	15 50	-46. 0 -32. 0 -32. 0 -18. 0	314. 3 328. 3 314. 9 328. 9	+9.0 +10.0 +9.0 +11.0	93 154 110 126		247 236
Jan. 8 (Naval Observatory)	13	1	-18. 0 -4. 0	315.6	+10.0 +11.0	123 154		277
Jan. 9 (Mount Wilson)	13	50	-5.0 + 9.0	315. 0 329. 0	$+9.0 \\ +11.0$	109 166		275
Jan. 10 (Naval Observatory)	11	55	$+8.0 \\ +22.0$	329.8	$+10.0 \\ +11.0$	139	154	293
Jan. 11 (Perkins Observatory)	15	40	$ +20.0 \\ +32.0$		$+11.0 \\ +11.0$	70 70		140

## POSITIONS AND AREAS OF SUN SPOTS-Continued

[Communicated by Capt. J. F. Hellweg, Superintendent United States Naval Observatory. Data furnished by Naval Observatory, in cooperation with Harvard, Perkins, and Mount Wilson Observatories. The differences of longitude are measured from central meridian, positive west. The north latitudes are plus. Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere. The total area, including spots and groups, is given for each day in the last column]

		Eastern		Heliographic			Area	
Date	standard civil time		Diff. long.	Longi- tude	Lati- tude	Spot	Group	area for each day
1933	h.	m.	۰	0	•			
Jan. 12 (Naval Observatory)	13	30	+4.0 +36.0	284. 6 316. 6	+8.0 +10.0	123	31	
Jan. 13 (Perkins Observatory)	15	0	$+49.0 \\ +5.0 \\ +34.0$	329. 6 271. 7 300. 7	+11.0 +5.0 +15.0	139	45 50	293
Jan. 14 (Mount Wilson)	12	0	+46.0 $+59.0$ $+20.0$ $+51.0$	312. 7 325. 7 275. 1 306. 1	+11.0 $+11.0$ $+3.0$ $+15.0$	70 70	53 74	235
Jan. 15 (Naval Observatory)	13	16	+62.0 +75.0 +33.0 +66.0	317.1	+10.0 $+11.0$ $+3.0$ $+14.0$	85 171	37 62	388
Jan. 16 (Naval Observatory)	10	43		317. 2 275. 5 No spots		108	19	207
Jan. 18 (Harvard Observatory) Jan. 19 (Perkins Observatory)	13	15		No spots				
Jan. 20 (Naval Observatory)		5		No spots				
Jan. 22 (Naval Observatory)			No spots.					
Jan. 23 (Naval Observatory)	10	20		No spots				
Jan. 24 (Naval Observatory)				No spots				
Jan. 28 (Naval Observatory)	10	47	-58.0				31	31
Jan. 29 (Naval Observatory)	11	19	-44.0	14.0	+5.0	100	154	154
Jan. 30 (Naval Observatory)	10	56	-79.0 $-33.0$	326. 0 12. 0	$+12.0 \\ +4.0$	123	50	
	İ		-26.0	19.0	+5.0		77	250
Jan. 31 (Naval Observatory)	11	5	-88.0	303. S	+16.0	247		
			-64.0 $-15.0$	327. 8 16. 8	+12.0 +6.0		108 216	571
Mean daily area for Janu-			-13.0	10.8	70.0		210	}
ary				.				17:

## PROVISIONAL SUN-SPOT RELATIVE NUMBERS FOR JANUARY, 1933

(Dependent alone on observations at Zurich and its station at Arosa)

[Data furnished through the courtesy of Prof. W. Brunner, University of Zurich, Switzerland]

January, 1933	Relative numbers	January, 1933	Relative numbers	January, 1933	Relative numbers
1	0 0 d 7 14 17 17 19 	11	17 Mc 26 37 32 27 14 7 0	21 22 23 24 25 26 27 28 29 30	0 0 0 0 0 0 0 Ec 9 10

Mean: 27 days=11.3.

a=Passage of an average-sized group through the central meridian. b=Passage of a large group or spot through the central meridian. c=New formation of a center of activity: E, on the eastern part of the sun's disk; W, on the western part; M, in the central zone. d=Entrance of a large or average-sized center of activity on the east limb.

## AEROLOGICAL OBSERVATIONS

[Aerological Division, W. R. Gregg, in charge]

By L. T. SAMUELS

Free-air temperatures during January averaged decidedly above normal except at San Diego, where small negative departures occurred. (See Table 1.) The largest positive departures occurred in the lower levels at Omaha and Washington. Relative humidities averaged close to normal at most stations, with the greatest deviations occurring at Omaha where the departures were negative.

In harmony with the large positive temperature departures, the monthly free-air resultant wind directions contained either greater southerly or smaller northerly components than normal at the corresponding stations. In most cases the resultant velocities were above normal. At 3,000 meters elevation and above, however, but little deviation from the normal directions occurred.

Airplane observations were made daily throughout the month at the Weather Bureau stations, with the exception of the 21st at Chicago when, dense fog continued all day.